

The Evolution of the Physicist's Picture of Nature

An account of how physical theory has developed in the past and how, in the light of this development, it can perhaps be expected to develop in the future

by P. A. M. Dirac

In this article I should like to discuss the development of general physical theory: how it developed in the past and how one may expect it to develop in the future. One can look on this continual development as a process of evolution, a process that has been going on for several centuries.

The first main step in this process of evolution was brought about by Newton. Before Newton, people looked on the world as being essentially two-dimensional—the two dimensions in which one can walk about—and the up-and-down dimension seemed to be something essentially different. Newton showed how one can look on the up-and-down direction as being symmetrical with the other two directions, by bringing in gravitational forces and showing how they take their place in physical theory. One can say that Newton enabled us to pass from a picture with two-dimensional symmetry to a picture with three-dimensional symmetry.

Einstein made another step in the same direction, showing how one can pass from a picture with three-dimensional symmetry to a picture with four-dimensional symmetry. Einstein brought in time and showed how it plays a role that is in many ways symmetrical with the three space dimensions. However, this symmetry is not quite perfect. With

Einstein's picture one is led to think of the world from a four-dimensional point of view, but the four dimensions are not completely symmetrical. There are some directions in the four-dimensional picture that are different from others: directions that are called null directions, along which a ray of light can move; hence the four-dimensional picture is not completely symmetrical. Still, there is a great deal of symmetry among the four dimensions. The only lack of symmetry, so far as concerns the equations of physics, is in the appearance of a minus sign in the equations with respect to the time dimension as compared with the three space dimensions [see top equation on page 50].

We have, then, the development from the three-dimensional picture of the world to the four-dimensional picture. The reader will probably not be happy with this situation, because the world still appears three-dimensional to his consciousness. How can one bring this appearance into the four-dimensional picture that Einstein requires the physicist to have?

What appears to our consciousness is really a three-dimensional section of the four-dimensional picture. We must take a three-dimensional section to give us what appears to our consciousness at one time; at a later time we shall have a

different three-dimensional section. The task of the physicist consists largely of relating events in one of these sections to events in another section referring to a later time. Thus the picture with four-dimensional symmetry does not give us the whole situation. This becomes particularly important when one takes into account the developments that have been brought about by quantum theory. Quantum theory has taught us that we have to take the process of observation into account, and observations usually require us to bring in the three-dimensional sections of the four-dimensional picture of the universe.

The special theory of relativity, which Einstein introduced, requires us to put all the laws of physics into a form that displays four-dimensional symmetry. But when we use these laws to get results about observations, we have to bring in something additional to the four-dimensional symmetry, namely the three-dimensional sections that describe our consciousness of the universe at a certain time.

Einstein made another most important contribution to the development of our physical picture: he put forward the general theory of relativity, which requires us to suppose that the space of physics is curved. Before this physicists

had always worked with a flat space, the three-dimensional flat space of Newton which was then extended to the four-dimensional flat space of special relativity. General relativity made a really important contribution to the evolution of our physical picture by requiring us to go over to curved space. The general requirements of this theory mean that all the laws of physics can be formulated in curved four-dimensional space, and that they show symmetry among the four dimensions. But again, when we want to bring in observations, as we must if we look at things from the point of view of quantum theory, we have to refer to a section of this four-dimensional space. With the four-dimensional space curved, any section that we make in it also has to be curved, because in general we cannot give a meaning to a flat section in a curved space. This leads us to a picture in which we have to take curved three-dimensional sections in the curved four-dimensional space and discuss observations in these sections.

During the past few years people have been trying to apply quantum ideas to

gravitation as well as to the other phenomena of physics, and this has led to a rather unexpected development, namely that when one looks at gravitational theory from the point of view of the sections, one finds that there are some degrees of freedom that drop out of the theory. The gravitational field is a tensor field with 10 components. One finds that six of the components are adequate for describing everything of physical importance and the other four can be dropped out of the equations. One cannot, however, pick out the six important components from the complete set of 10 in any way that does not destroy the four-dimensional symmetry. Thus if one insists on preserving four-dimensional symmetry in the equations, one cannot adapt the theory of gravitation to a discussion of measurements in the way quantum theory requires without being forced to a more complicated description than is needed by the physical situation. This result has led me to doubt how fundamental the four-dimensional requirement in physics is. A few decades ago it seemed quite certain that one had


to express the whole of physics in four-dimensional form. But now it seems that four-dimensional symmetry is not of such overriding importance, since the description of nature sometimes gets simplified when one departs from it.

Now I should like to proceed to the developments that have been brought about by quantum theory. Quantum theory is the discussion of very small things, and it has formed the main subject of physics for the past 60 years. During this period physicists have been amassing quite a lot of experimental information and developing a theory to correspond to it, and this combination of theory and experiment has led to important developments in the physicist's picture of the world.

The quantum first made its appearance when Planck discovered the need to suppose that the energy of electromagnetic waves can exist only in multiples of a certain unit, depending on the frequency of the waves, in order to explain the law of black-body radiation. Then Einstein discovered the same unit of energy occurring in the photoelectric effect. In this early work on quantum theory one simply had to accept the unit of energy without being able to incorporate it into a physical picture.

The first new picture that appeared was Bohr's picture of the atom. It was a picture in which we had electrons moving about in certain well-defined orbits and occasionally making a jump from one orbit to another. We could not picture how the jump took place. We just had to accept it as a kind of discontinuity. Bohr's picture of the atom worked only for special examples, essentially when there was only one electron that was of importance for the problem under consideration. Thus the picture was an incomplete and primitive one.

The big advance in the quantum theory came in 1925, with the discovery of quantum mechanics. This advance was brought about independently by two men, Heisenberg first and Schrödinger soon afterward, working from different points of view. Heisenberg worked keeping close to the experimental evidence about spectra that was being amassed at that time, and he found out how the experimental information could be fitted into a scheme that is now known as matrix mechanics. All the experimental data of spectroscopy fitted beautifully into the scheme of matrix mechanics, and this led to quite a different picture of the atomic world. Schrödinger worked from a more mathematical point of view, trying to find a beautiful theory for describ-



ISAAC NEWTON (1642-1727), with his law of gravitation, changed the physicist's picture of nature from one with two-dimensional symmetry to one with three-dimensional symmetry. This drawing of him was made in 1760 by James Macardel from a painting by Enoch Seeman.

ing atomic events, and was helped by De Broglie's ideas of waves associated with particles. He was able to extend De Broglie's ideas and to get a very beautiful equation, known as Schrödinger's wave equation, for describing atomic processes. Schrödinger got this equation by pure thought, looking for some beautiful generalization of De Broglie's ideas, and not by keeping close to the experimental development of the subject in the way Heisenberg did.

I might tell you the story I heard from Schrödinger of how, when he first got the idea for this equation, he immediately applied it to the behavior of the electron in the hydrogen atom, and then he got results that did not agree with experiment. The disagreement arose because at that time it was not known that the electron has a spin. That, of course, was a great disappointment to Schrödinger, and it caused him to abandon the work for some months. Then he noticed that if he applied the theory in a more approximate way, not taking into account the refinements required by relativity, to this rough approximation his work was in agreement with observation. He published his first paper with only this rough approximation, and in that way Schrödinger's wave equation was presented to the world. Afterward, of course, when people found out how to take into account correctly the spin of the electron, the discrepancy between the results of applying Schrödinger's relativistic equation and the experiments was completely cleared up.

I think there is a moral to this story, namely that it is more important to have beauty in one's equations than to have them fit experiment. If Schrödinger had been more confident of his work, he could have published it some months earlier, and he could have published a more accurate equation. That equation is now known as the Klein-Gordon equation, although it was really discovered by Schrödinger, and in fact was discovered by Schrödinger before he discovered his nonrelativistic treatment of the hydrogen atom. It seems that if one is working from the point of view of getting beauty in one's equations, and if one has really a sound insight, one is on a sure line of progress. If there is not complete agreement between the results of one's work and experiment, one should not allow oneself to be too discouraged, because the discrepancy may well be due to minor features that are not properly taken into account and that will get cleared up with further developments of the theory.



ALBERT EINSTEIN (1879–1955), with his special theory of relativity, changed the physicist's picture of nature from one with three-dimensional symmetry to one with four-dimensional symmetry. This photograph of him and his wife and their daughter Margot was made in 1929.


That is how quantum mechanics was discovered. It led to a drastic change in the physicist's picture of the world, perhaps the biggest that has yet taken place. This change comes from our having to give up the deterministic picture we had always taken for granted. We are led to a theory that does not predict with certainty what is going to happen in the future but gives us information only about the probability of occurrence of various events. This giving up of determinacy has been a very controversial subject, and some people do not like it at all. Einstein in particular never liked it.

Although Einstein was one of the great contributors to the development of quantum mechanics, he still was always rather hostile to the form that quantum mechanics evolved into during his lifetime and that it still retains.

The hostility some people have to the giving up of the deterministic picture can be centered on a much discussed paper by Einstein, Podolsky and Rosen dealing with the difficulty one has in forming a consistent picture that still gives results according to the rules of quantum mechanics. The rules of quantum mechanics are quite definite. People



NIELS BOHR (1885–1962) introduced the idea that the electron moved about the nucleus in well-defined orbits. This photograph was made in 1922, nine years after the publication of his paper.



MAX PLANCK (1858–1947) introduced the idea that electromagnetic radiation consists of quanta, or particles. This photograph was made in 1913, 13 years after his original paper was published.

know how to calculate results and how to compare the results of their calculations with experiment. Everyone is agreed on the formalism. It works so well that nobody can afford to disagree with it. But still the picture that we are to set up behind this formalism is a subject of controversy.

I should like to suggest that one not worry too much about this controversy. I feel very strongly that the stage physics has reached at the present day is not the final stage. It is just one stage in the evolution of our picture of nature, and we should expect this process of evolution to continue in the future, as biological evolution continues into the future. The present stage of physical theory is merely a steppingstone toward the better stages we shall have in the future. One can be quite sure that there will be better stages simply because of the difficulties that occur in the physics of today.

I should now like to dwell a bit on the difficulties in the physics of the present day. The reader who is not an expert in the subject might get the idea that because of all these difficulties physical theory is in pretty poor shape and that the quantum theory is not much good. I should like to correct this impression by saying that quantum theory is an extremely good theory. It gives wonderful agreement with observation over a wide range of phenomena. There is no doubt that it is a good theory, and the only reason physicists talk so much about

the difficulties in it is that it is precisely the difficulties that are interesting. The successes of the theory are all taken for granted. One does not get anywhere simply by going over the successes again and again, whereas by talking over the difficulties people can hope to make some progress.

The difficulties in quantum theory are of two kinds. I might call them Class One difficulties and Class Two difficulties. Class One difficulties are the difficulties I have already mentioned: How can one form a consistent picture behind the rules for the present quantum theory? These Class One difficulties do not really worry the physicist. If the physicist knows how to calculate results and compare them with experiment, he is quite happy if the results agree with his experiments, and that is all he needs. It is only the philosopher, wanting to have a satisfying description of nature, who is bothered by Class One difficulties.

There are, in addition to the Class One difficulties, the Class Two difficulties, which stem from the fact that the present laws of quantum theory are not always adequate to give any results. If one pushes the laws to extreme conditions—to phenomena involving very high energies or very small distances—one sometimes gets results that are ambiguous or not really sensible at all. Then it is clear that one has reached the limits of application of the theory and that some further development is needed. The Class Two difficulties are important even for

the physicist, because they put a limitation on how far he can use the rules of quantum theory to get results comparable with experiment.

I should like to say a little more about the Class One difficulties. I feel that one should not be bothered with them too much, because they are difficulties that refer to the present stage in the development of our physical picture and are almost certain to change with future development. There is one strong reason, I think, why one can be quite confident that these difficulties will change. There are some fundamental constants in nature: the charge on the electron (designated e), Planck's constant divided by 2π (designated \hbar) and the velocity of light (c). From these fundamental constants one can construct a number that has no dimensions: the number $\hbar c/e^2$. That number is found by experiment to have the value 137, or something very close to 137. Now, there is no known reason why it should have this value rather than some other number. Various people have put forward ideas about it, but there is no accepted theory. Still, one can be fairly sure that someday physicists will solve the problem and explain why the number has this value. There will be a physics in the future that works when $\hbar c/e^2$ has the value 137 and that will not work when it has any other value.

The physics of the future, of course, cannot have the three quantities \hbar , e and c all as fundamental quantities. Only two

of them can be fundamental, and the third must be derived from those two. It is almost certain that c will be one of the two fundamental ones. The velocity of light, c , is so important in the four-dimensional picture, and it plays such a fundamental role in the special theory of relativity, correlating our units of space and time, that it has to be fundamental. Then we are faced with the fact that of the two quantities \hbar and e , one will be fundamental and one will be derived. If \hbar is fundamental, e will have to be explained in some way in terms of the square root of \hbar , and it seems most unlikely that any fundamental theory can give e in terms of a square root, since square roots do not occur in basic equations. It is much more likely that e will be the fundamental quantity and that \hbar will be explained in terms of e^2 . Then there will be no square root in the basic equations. I think one is on safe ground if one makes the guess that in the physical picture we shall have at some future stage e and c will be fundamental quantities and \hbar will be derived.

If \hbar is a derived quantity instead of a fundamental one, our whole set of ideas about uncertainty will be altered: \hbar is the fundamental quantity that occurs in the Heisenberg uncertainty relation connecting the amount of uncertainty in a position and in a momentum. This uncertainty relation cannot play a fundamental role in a theory in which \hbar itself is not a fundamental quantity. I think one can make a safe guess that uncertainty relations in their present form will not survive in the physics of the future.


Of course there will not be a return to the determinism of classical physical theory. Evolution does not go backward. It will have to go forward. There will have to be some new development that is quite unexpected, that we cannot make a guess about, which will take us still further from classical ideas but which will alter completely the discussion of uncertainty relations. And when this new development occurs, people will find it all rather futile to have had so much of a discussion on the role of observation in the theory, because they will have then a much better point of view from which to look at things. So I shall say that if we can find a way to describe the uncertainty relations and the indeterminacy of present quantum mechanics that is satisfying to our philosophical ideas, we can count ourselves lucky. But if we cannot find such a way, it is nothing to be really disturbed about. We simply have to take into account that we are at a transitional stage

and that perhaps it is quite impossible to get a satisfactory picture for this stage.

I have disposed of the Class One difficulties by saying that they are really not so important, that if one can make progress with them one can count oneself lucky, and that if one cannot it is nothing to be genuinely disturbed about. The Class Two difficulties are the really serious ones. They arise primarily from the fact that when we apply our quantum theory to fields in the way we have to if we are to make it agree with special relativity, interpreting it in terms of the three-dimensional sections I have mentioned, we have equations that at first look all right. But when one tries to solve them, one finds that they do not have any solutions. At this point we ought to say that we do not have a theory. But physicists are very ingenious about it, and they have found a way to make progress in spite of this obstacle. They find that when they try to solve the equations, the trouble is that certain quantities that ought to be finite are actually infinite. One gets integrals that diverge instead of converging to something definite. Physicists have found that there is a

way to handle these infinities according to certain rules, which makes it possible to get definite results. This method is known as the renormalization method.

I shall merely explain the idea in words. We start out with a theory involving equations. In these equations there occur certain parameters: the charge of the electron, e , the mass of the electron, m , and things of a similar nature. One then finds that these quantities, which appear in the original equations, are not equal to the measured values of the charge and the mass of the electron. The measured values differ from these by certain correcting terms— Δe , Δm and so on—so that the total charge is $e + \Delta e$ and the total mass $m + \Delta m$. These changes in charge and mass are brought about through the interaction of our elementary particle with other things. Then one says that $e + \Delta e$ and $m + \Delta m$, being the observed things, are the important things. The original e and m are just mathematical parameters; they are unobservable and therefore just tools one can discard when one has got far enough to bring in the things that one can com-



LOUIS DE BROGLIE (1892–) put forward the idea that particles are associated with waves. This photograph was made in 1929, five years after the appearance of his paper.

pare with observation. This would be a quite correct way to proceed if Δe and Δm were small (or even if they were not so small but finite) corrections. According to the actual theory, however, Δe and Δm are infinitely great. In spite of that fact one can still use the formalism and get results in terms of $e + \Delta e$ and $m + \Delta m$, which one can interpret by saying that the original e and m have to be minus infinity of a suitable amount to compensate for the Δe and Δm that are infinitely great. One can use the theory to get results that can be compared with experiment, in particular for electrodynamics. The surprising thing is that in the case of electrodynamics one gets results that are in extremely good agreement with experiment. The agreement applies to many significant figures—the kind of accuracy that previously one had only in astronomy. It is because of this good agreement that physicists do attach some value to the renormalization theory, in spite of its illogical character.

It seems to be quite impossible to put this theory on a mathematically sound basis. At one time physical theory was all built on mathematics that was inherently

sound. I do not say that physicists always use sound mathematics; they often use unsound steps in their calculations. But previously when they did so it was simply because of, one might say, laziness. They wanted to get results as quickly as possible without doing unnecessary work. It was always possible for the pure mathematician to come along and make the theory sound by bringing in further steps, and perhaps by introducing quite a lot of cumbersome notation and other things that are desirable from a mathematical point of view in order to get everything expressed rigorously but do not contribute to the physical ideas. The earlier mathematics could always be made sound in that way, but in the renormalization theory we have a theory that has defied all the attempts of the mathematician to make it sound. I am inclined to suspect that the renormalization theory is something that will not survive in the future, and that the remarkable agreement between its results and experiment should be looked on as a fluke.

This is perhaps not altogether surprising, because there have been similar flukes in the past. In fact, Bohr's elec-

tron-orbit theory was found to give very good agreement with observation as long as one confined oneself to one-electron problems. I think people will now say that this agreement was a fluke, because the basic ideas of Bohr's orbit theory have been superseded by something radically different. I believe the successes of the renormalization theory will be on the same footing as the successes of the Bohr orbit theory applied to one-electron problems.

The renormalization theory has removed some of these Class Two difficulties, if one can accept the illogical character of discarding infinities, but it does not remove all of them. There are a good many problems left over concerning particles other than those that come into electrodynamics: the new particles—mesons of various kinds and neutrinos. There the theory is still in a primitive stage. It is fairly certain that there will have to be drastic changes in our fundamental ideas before these problems can be solved.

One of the problems is the one I have already mentioned about accounting for the number 137. Other problems are how to introduce the fundamental length to physics in some natural way, how to explain the ratios of the masses of the elementary particles and how to explain their other properties. I believe separate ideas will be needed to solve these distinct problems and that they will be solved one at a time through successive stages in the future evolution of physics. At this point I find myself in disagreement with most physicists. They are inclined to think one master idea will be discovered that will solve all these problems together. I think it is asking too much to hope that anyone will be able to solve all these problems together. One should separate them one from another as much as possible and try to tackle them separately. And I believe the future development of physics will consist of solving them one at a time, and that after any one of them has been solved there will still be a great mystery about how to attack further ones.

I might perhaps discuss some ideas I have had about how one can possibly attack some of these problems. None of these ideas has been worked out very far, and I do not have much hope for any one of them. But I think they are worth mentioning briefly.

One of these ideas is to introduce something corresponding to the luminiferous ether, which was so popular among the physicists of the 19th century. I said earlier that physics does not evolve back-

$$ds^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2$$

FOUR-DIMENSIONAL SYMMETRY introduced by the special theory of relativity is not quite perfect. This equation is the expression for the invariant distance in four-dimensional space-time. The symbol s is the invariant distance; c , the speed of light; t , time; x , y and z , the three spatial dimensions. The d 's are differentials. The lack of complete symmetry lies in the fact that the contribution from the time direction ($c^2 dt^2$) does not have the same sign as the contributions from the three spatial directions ($-dx^2$, $-dy^2$ and $-dz^2$).

$$\left(\frac{i\hbar}{2\pi c} \frac{\partial}{\partial t} + \frac{e^2}{cr} \right) \psi = \left[m^2 c^2 - \frac{\hbar^2}{4\pi^2} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \right] \psi$$

SCHRÖDINGER'S FIRST WAVE EQUATION did not fit experimental results because it did not take into account the spin of the electron, which was not known at the time. The equation is a generalization of De Broglie's equation for the motion of a free electron. The symbol e represents the charge on the electron; i , the square root of minus one; \hbar , Planck's constant; r , the distance from the nucleus; ψ , Schrödinger's wave function; m , the mass of the electron. The symbols resembling sixes turned backward are partial derivatives.

$$\left(E + \frac{e^2}{r} \right) \psi = - \frac{\hbar^2}{8\pi^2 m} \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \psi$$

SCHRÖDINGER'S SECOND WAVE EQUATION is an approximation to the original equation, which does not take into account the refinements that are required by relativity.

ward. When I talk about reintroducing the ether, I do not mean to go back to the picture of the ether that one had in the 19th century, but I do mean to introduce a new picture of the ether that will conform to our present ideas of quantum theory. The objection to the old idea of the ether was that if you suppose it to be a fluid filling up the whole of space, in any place it has a definite velocity, which destroys the four-dimensional symmetry required by Einstein's special principle of relativity. Einstein's special relativity killed this idea of the ether.

But with our present quantum theory we no longer have to attach a definite velocity to any given physical thing, because the velocity is subject to uncertainty relations. The smaller the mass of the thing we are interested in, the more important are the uncertainty relations. Now, the ether will certainly have very little mass, so that uncertainty relations for it will be extremely important. The velocity of the ether at some particular place should therefore not be pictured as definite, because it will be subject to uncertainty relations and so may be anything over a wide range of values. In that way one can get over the difficulties of reconciling the existence of an ether with the special theory of relativity.

There is one important change this will make in our picture of a vacuum. We would like to think of a vacuum as a region in which we have complete symmetry between the four dimensions of space-time as required by special relativity. If there is an ether subject to uncertainty relations, it will not be possible to have this symmetry accurately. We can suppose that the velocity of the ether is equally likely to be anything within a wide range of values that would give the symmetry only approximately. We cannot in any precise way proceed to the limit of allowing all values for the velocity between plus and minus the velocity of light, which we would have to do in order to make the symmetry accurate. Thus the vacuum becomes a state that is unattainable. I do not think that this is a physical objection to the theory. It would mean that the vacuum is a state we can approach very closely. There is no limit as to how closely we can approach it, but we can never attain it. I believe that would be quite satisfactory to the experimental physicist. It would, however, mean a departure from the notion of the vacuum that we have in the quantum theory, where we start off with the vacuum state having exactly the symmetry required by special relativity.

That is one idea for the development of physics in the future that would



ERWIN SCHRÖDINGER (1887–1961) devised his wave equation by extending De Broglie's idea that waves are associated with particles to the electrons moving around the nucleus. This photograph was made in 1929, four years after he had published his second equation.

change our picture of the vacuum, but change it in a way that is not unacceptable to the experimental physicist. It has proved difficult to continue with the theory, because one would need to set up mathematically the uncertainty relations for the ether and so far some satisfactory theory along these lines has not been discovered. If it could be developed satisfactorily, it would give rise to a new kind of field in physical theory, which might help in explaining some of the elementary particles.

Another possible picture I should like to mention concerns the question of why all the electric charges that are observed in nature should be multiples of one elementary unit, e . Why does one not have a continuous distribution of charge occurring in nature? The picture I propose goes back to the idea of Faraday lines of force and involves a development of this idea. The Faraday

lines of force are a way of picturing electric fields. If we have an electric field in any region of space, then according to Faraday we can draw a set of lines that have the direction of the electric field. The closeness of the lines to one another gives a measure of the strength of the field—they are close where the field is strong and less close where the field is weak. The Faraday lines of force give us a good picture of the electric field in classical theory.

When we go over to quantum theory, we bring a kind of discreteness into our basic picture. We can suppose that the continuous distribution of Faraday lines of force that we have in the classical picture is replaced by just a few discrete lines of force with no lines of force between them.

Now, the lines of force in the Faraday picture end where there are charges. Therefore with these quantized Faraday lines of force it would be reasonable to

suppose the charge associated with each line, which has to lie at the end if the line of force has an end, is always the same (apart from its sign), and is always just the electronic charge, $-e$ or $+e$. This leads us to a picture of discrete Faraday lines of force, each associated with a charge, $-e$ or $+e$. There is a direction attached to each line, so that the ends of a line that has two ends are not the same, and there is a charge $+e$ at one end and a charge $-e$ at the other. We may have lines of force extending to infinity, of course, and then there is no charge.

If we suppose that these discrete Faraday lines of force are something basic in physics and lie at the bottom of our picture of the electromagnetic field, we shall have an explanation of why charges always occur in multiples of e . This happens because if we have any particle with some lines of force ending on it, the number of these lines must be a whole number. In that way we get a picture that is qualitatively quite reasonable.

We suppose these lines of force can

move about. Some of them, forming closed loops or simply extending from minus infinity to infinity, will correspond to electromagnetic waves. Others will have ends, and the ends of these lines will be the charges. We may have a line of force sometimes breaking. When that happens, we have two ends appearing, and there must be charges at the two ends. This process—the breaking of a line of force—would be the picture for the creation of an electron (e^-) and a positron (e^+). It would be quite a reasonable picture, and if one could develop it, it would provide a theory in which e appears as a basic quantity. I have not yet found any reasonable system of equations of motion for these lines of force, and so I just put forward the idea as a possible physical picture we might have in the future.

There is one very attractive feature in this picture. It will quite alter the discussion of renormalization. The renormalization we have in our present quantum electrodynamics comes from starting off with what people call a bare electron—an electron without a charge

on it. At a certain stage in the theory one brings in the charge and puts it on the electron, thereby making the electron interact with the electromagnetic field. This brings a perturbation into the equations and causes a change in the mass of the electron, the Δm , which is to be added to the previous mass of the electron. The procedure is rather roundabout because it starts off with the unphysical concept of the bare electron. Probably in the improved physical picture we shall have in the future the bare electron will not exist at all.

Now, that state of affairs is just what we have with the discrete lines of force. We can picture the lines of force as strings, and then the electron in the picture is the end of a string. The string itself is the Coulomb force around the electron. A bare electron means an electron without the Coulomb force around it. That is inconceivable with this picture, just as it is inconceivable to think of the end of a piece of string without thinking of the string itself. This, I think, is the kind of way in which we should try to develop our physical picture—to bring in ideas that make inconceivable the things we do not want to have. Again we have a picture that looks reasonable, but I have not found the proper equations for developing it.

I might mention a third picture with which I have been dealing lately. It involves departing from the picture of the electron as a point and thinking of it as a kind of sphere with a finite size. Of course, it is really quite an old idea to picture the electron as a sphere, but previously one had the difficulty of discussing a sphere that is subject to acceleration and to irregular motion. It will get distorted, and how is one to deal with the distortions? I propose that one should allow the electron to have, in general, an arbitrary shape and size. There will be some shapes and sizes in which it has less energy than in others, and it will tend to assume a spherical shape with a certain size in which the electron has the least energy.

This picture of the extended electron has been stimulated by the discovery of the mu meson, or muon, one of the new particles of physics. The muon has the surprising property of being almost identical with the electron except in one particular, namely, its mass is some 200 times greater than the mass of the electron. Apart from this disparity in mass the muon is remarkably similar to the electron, having, to an extremely high degree of accuracy, the same spin and the same magnetic moment in proportion to its mass as the electron does. This




WERNER HEISENBERG (1901–) introduced matrix mechanics, which, like the Schrödinger theory, accounted for the motions of the electron. This photograph was made in 1929.

leads to the suggestion that the muon should be looked on as an excited electron. If the electron is a point, picturing how it can be excited becomes quite awkward. But if the electron is the most stable state for an object of finite size, the muon might just be the next most stable state in which the object undergoes a kind of oscillation. That is an idea I have been working on recently. There are difficulties in the development of this idea, in particular the difficulty of bringing in the correct spin.

I have mentioned three possible ways in which one might think of developing our physical picture. No doubt there will be others that other people will think of. One hopes that sooner or later someone will find an idea that really fits and leads to a big development. I am rather pessimistic about it and am inclined to think none of them will be good enough. The future evolution of basic physics—that is to say, a development that will really solve one of the fundamental problems, such as bringing in the fundamental length or calculating the ratio of the masses—may require some much more drastic change in our physical picture. This would mean that in our present attempts to think of a new physical picture we are setting our imaginations to work in terms of inadequate physical concepts. If that is really the case, how can we hope to make progress in the future?

There is one other line along which one can still proceed by theoretical means. It seems to be one of the fundamental features of nature that fundamental physical laws are described in terms of a mathematical theory of great beauty and power, needing quite a high standard of mathematics for one to understand it. You may wonder: Why is nature constructed along these lines? One can only answer that our present knowledge seems to show that nature is so constructed. We simply have to accept it. One could perhaps describe the situation by saying that God is a mathematician of a very high order, and He used very advanced mathematics in constructing the universe. Our feeble attempts at mathematics enable us to understand a bit of the universe, and as we proceed to develop higher and higher mathematics we can hope to understand the universe better.

This view provides us with another way in which we can hope to make advances in our theories. Just by studying mathematics we can hope to make a guess at the kind of mathematics that will come into the physics of the future.



LINES OF FORCE in an electromagnetic field, if they are assumed to be discrete in the quantum theory, suggest why electric charges always occur in multiples of the charge of the electron. In Dirac's view, when a line of force has two ends, there is a particle with charge $-e$, perhaps an electron, at one end and a particle with charge $+e$, perhaps a positron, at the other end. When a closed line of force is broken, an electron-positron pair materializes.

A good many people are working on the mathematical basis of quantum theory, trying to understand the theory better and to make it more powerful and more beautiful. If someone can hit on the right lines along which to make this development, it may lead to a future advance in which people will first discover the equations and then, after examining them, gradually learn how to apply them. To some extent that corresponds with the line of development that occurred with Schrödinger's discovery of his wave equation. Schrödinger discovered the equation simply by looking for an equation with mathematical beauty. When the equation was first discovered, people saw that it fitted in certain ways, but the general principles according to which one should apply it were worked out only some two or three years later. It may well be that the next advance in physics will come about along these lines: people first discovering the equa-

tions and then needing a few years of development in order to find the physical ideas behind the equations. My own belief is that this is a more likely line of progress than trying to guess at physical pictures.

Of course, it may be that even this line of progress will fail, and then the only line left is the experimental one. Experimental physicists are continuing their work quite independently of theory, collecting a vast storehouse of information. Sooner or later there will be a new Heisenberg who will be able to pick out the important features of this information and see how to use them in a way similar to that in which Heisenberg used the experimental knowledge of spectra to build his matrix mechanics. It is inevitable that physics will develop ultimately along these lines, but we may have to wait quite a long time if people do not get bright ideas for developing the theoretical side.